

The q -nonadditivity of nonextensive statistics is not a true physical property

Q.A. Wang[†], C.J. Ou[‡], A. J.C. Chen[§]

[†]*ISMANS, 44, Avenue F.A. Bartholdi, Le Mans, France*

[‡]*College of Information Science and Engineering,
Huaqiao University, Quanzhou, China*

[§]*Department of Physics, Xiamen University, Xiamen, China*

Abstract

This is a note showing that, contrary to our lasting belief, the nonadditivity $X(1 \cup 2) = X(1) + X(2) + \alpha X(1)X(2)$ is not a true physical property. α in this expression cannot be unique for a given system. It unavoidably depends on how one mathematically divides the system and cannot be used to characterize nonadditivity. As a matter of fact, its use is mathematically inconsistent.

People who worked with q -nonextensive statistics[1] are all aware of a nonlinear relationship characterizing the nonadditivity of entropy and energy which is an emblem of the theory and a central object of polemics about the foundation of the theory (see for example [2] and references there-in) and its relationship with thermodynamics (a brief review can be found in [3]). The nonadditivity is given by, for a total system composed of two subsystems 1 and 2:

$$X(1 \cup 2) = X(1) + X(2) + \alpha X(1)X(2) \quad (1)$$

where X represents entropy, energy or other variables, $\alpha = 1 - q$ (or $q - 1$) (with positive q) is a real coupling constant between 1 and 2. For the total system, as claimed in every statement relative to this equality, α characterizes its nonadditivity. Hence for a given system and situation, α is expected to

be unique. This relationship has been widely employed in the definition of temperature, pressure and other intensive variables in order to relate the nonextensive statistical mechanics to thermodynamics.

The origine of this nonadditivity was the reference [4] from the hypothesis of product join probability $p(1+2) = p(1)p(2)$ in order to show the nonadditivity of entropy S given by $S(1 \cup 2) = S(1) + S(2) + \alpha S(1)S(2)$ which was subsequently used in the discussion of zeroth law of thermodynamics with an additional nonadditivity of energy U given by $U(1 \cup 2) = U(1) + U(2) + \alpha U(1)U(2)$ (see for example a historical review in [3]).

We note here that this nonadditivity is false because, on the one hand, α is necessarily dependent on the subsystems of a given system so that it cannot be used for characterizing the (unique) nonadditivity of the latter and, on the other hand, with α dependent on the subsystems, a self-contradiction is unavoidable when one treats a system composed of more than two subsystems.

Let us see a calculation for (homogeneous) gravitational systems[5] which shows that $\alpha_{1,2}$ indeed depends on the size of the two subsystems. The gravitational potential energy V of a solid spherical mass of radius r_2 separated into a smaller inner sphere of radius $r_1 < r_2$ and a spherical outer shell of thickness $r_2 - r_1$ is non-extensive: $V_{total} = V(1) + V(2) + \alpha_{1,2}V(1)V(2)$ where $\alpha_{1,2} = (\frac{r_1^3 r_2^2}{2} - \frac{r_1^5}{2}) / (\frac{r_1^5 r_2^5}{25} - \frac{r_1^8 r_2^2}{10} + \frac{3r_1^{10}}{50})$. It is obvious that this (homogeneous) system does not have unique α . Each way of partitioning the system will result in a different value of $\alpha_{1,2}$ or q , meaning that Eq.(1) does not have any practical meaning for characterizing the gravitation nonextensivity. It should be noticed that here we look at only the potential energy. If we add the thermal energy (heat) and the eventual kinetic energy of the global motion which are normally extensive and additive, the nonadditivity of the total energy cannot even take the form of Eq.(1).

Then a question arises: can Eq.(1) existe for certain if not all homogeneous system with α independent of the subsystems? Let us give an answer with entropy. For a given system divided into two subsystems 1 and 2, the total entropy is $S(1 \cup 2) = S(1) + S(2) + \alpha S(1)S(2)$. We can write $S(1)/S(2) = k_1/k_2$ where the positive ratio k_1/k_2 characterizes the division. This means that we can also write $S(1) = k_1 x$ and $S(2) = k_2 x$ here x is the common factor of $S(1)$ and $S(2)$. As a result, the total entropy reads $S(1 \cup 2) = (k_1 + k_2)x + \alpha k_1 k_2 x^2$. Since $S(1 \cup 2)$ is constant for a given system, we can let it be unity without lose of generality. This means

$(k_1 + k_2)x + \alpha k_1 k_2 x^2 = 1$. This equation must have positive solution for x (entropy is positive). That is $\Delta = (k_1 + k_2)^2 + 4\alpha k_1 k_2 \geq 0$ and $\alpha < \frac{\sqrt{\Delta} - (k_1 + k_2)}{2k_1 k_2}$. This last relationship shows that α cannot be independent of k_1 and k_2 , i.e., of the decomposition.

One can reach a more precise result simply by saying that, since the total system is given, Eq.(1) must read $X(1) + X(2) + \alpha X(1)X(2) = C$ where C is a constant. This implies $\alpha = \frac{C}{X(1)X(2)} - \frac{1}{X(2)} - \frac{1}{X(1)}$ so that α is in general dependent on $X(1)$ and $X(2)$. Unavoidably, each way of partitioning a given system yields a different value of α .

Now let us see a mathematical self-contradiction which happens in the application of Eq.(1). Suppose a total system partitioned into three subsystems 1, 2 and 3, which is a frequently encountered case in physics and chemistry. One can for example image three atomic clusters forming a bigger one, or the above mass sphere partitioned into an inner sphere and two exterior shells. Let the nonadditivity between the subsystems i and j be denoted by $X(i \cup j) = X(i) + X(j) + \alpha_{i,j}X(i)X(j)$ $i \neq j = 1, 2 \text{ or } 3$ where the constant $\alpha_{i,j}$ is the coupling constant of the composite system $i + j$, i.e., between the subsystems i and j . All three $\alpha_{i,j}$ can be different. Now let us see the already constructed total system with a given total $X(1 \cup 2 \cup 3)$ and imagine to add $X(1)$, $X(2)$ and $X(3)$ in different order. We first add $X(1)$ and $X(2)$ and then $X(3)$. The total $X(1 \cup 2 \cup 3)$ is given by

$$X((1 \cup 2) \cup 3) = X(1 \cup 2) + X(3) + \alpha_{(1,2),3}X(1 \cup 2)X(3) \quad (2)$$

$$\begin{aligned} &= X(1) + X(2) + X(3) + \alpha_{1,2}X(1)X(2) \\ &+ \alpha_{(1,2),3}[X(1)X(3) + X(2)X(3)] + \alpha_{1,2}\alpha_{(1,2),3}X(1)X(2)X(3) \end{aligned} \quad (3)$$

where $\alpha_{(1,2),3}$ is the coupling constant between the composite system $1 \cup 2$ and the subsystem 3. However, if we first add $X(1)$ and $X(3)$ and then $X(2)$, the result is

$$X((1 \cup 3) \cup 2) = X(1) + X(2) + X(3) + \alpha_{1,3}X(1)X(3) \quad (4)$$

$$+ \alpha_{(1,3),2}[X(1)X(2) + X(3)X(2)] + \alpha_{1,3}\alpha_{(1,3),2}X(1)X(2)X(3)$$

The above two operations must give the same result $X(1 \cup 2 \cup 3)$ since the additions of different order are just mathematical operations or thought experiences without physically disturbing the system. However, Eq.(2) can be equivalent to Eq.(4) only if all the coupling constants involved are zero or equal. The first case (zero coupling) is for extensive system. The second case is impossible as shown above.

The above analysis is simply mathematics. No doubt seems possible. One is obliged to say that Eq.(1) has no practical meaning in the description of nonextensive system. It must be rejected, together with the hypothesis of product join probability, from the nonextensive physics. The hypothetical product join probability, being the origin of the nonadditivity Eq.(1) for entropy and energy, is in fact unrealistic as a general rule for two subsystems which are dependent on each other due to nonextensivity. This hypothesis is in addition contradictory with probability theory. Although one can say that there may be dependent subsystems having by chance the product joint probability, such an accidental and unpredictable case must not be used for formulating general theory.

In summary, we showed that the nonadditivity given by Eq.(1) with constant α is untrue. Its use may lead to contradiction in the calculation of physical quantities for a composite system. We would like to emphasize that this unexpected result and its mathematical logic do not touch (as far as we see) the validity of the q -exponential probability distribution and the nonextensive statistics based on the Tsallis entropy as an axiom. However, since Eq.(1) was the starting point for the definition of intensive variables of nonextensive system, the already established (though discussible) connection between the nonextensive statistical mechanics and thermodynamics is definitely broken.

This is not an encouraging result, especially for the hard searching for the thermodynamics describing the systems that do not satisfy the conventional paradigms such as Gibbs-Shannon entropy formula, equilibrium condition, thermodynamic limits, ergodicity, energy and entropy additivity and others. Examples of these systems include finite system, long range interacting systems, large fluctuating system and many complex systems such as scale free network and social process to cite only some.

We thank Professor S. Abe for fruitful discussion.

References

- [1] C. Tsallis, Introduction to Nonextensive Statistical Mechanics, Springer 2009
- [2] Special issue of *Comptes Rendus Physique* **7**,(2006)
Special issue of *Europhysics news*, **36/6** (2005)

- [3] Q.A. Wang, Laurent Nivanen, Alain Le Mehaute, Michel Pezeril, *Europhys. Lett.*, **65**(2004)606
- [4] C. Tsallis, *J. Stat. Phys.* **52**, 479(1988)
- [5] C.J. Ou, et al, *Physica A* **387**, 5761(2008)